

A Northern Nevada Homeowner's Guide to Identifying and Managing Flixweed

Susan Donaldson, Water Quality and Weed Specialist

Wendy Hanson Mazet, Master Gardener Program Coordinator and Horticulturist

Other common names: Tansy mustard, herb sophia

Scientific name: *Descurainia sophia*

Family: Brassicaceae

Description: A bushy, much-branched plant that grows up to 2 or more feet tall, flixweed blooms early in the spring.

Leaves: The leaves are finely divided and hairy. The hairs are branched.

Stems: Stems are upright and branched. Plants grow in a rosette (ground-hugging form, see photo below right) until the flowering stems start growing.

Flowers: Tiny and yellow with four petals; arranged in branched structures. Blooms from early spring to summer.

Seeds: Produces narrow seed pods $\frac{1}{2}$ to $1\frac{1}{4}$ inches long.


Roots: Has a short taproot.

Native to: Europe; naturalized in much of the United States

Where it grows: Gardens, landscaped areas, rangeland, vacant lots, roadsides and other disturbed or unmanaged sites

Life cycle: Winter annual (sprouts in fall or early winter), summer annual (sprouts in spring or summer), sometimes biennial (flowers and dies in the second year of growth)

Reproduction: Reproduces by seed



Typical plant growing in disturbed site.



Rosettes have finely divided leaves.

Control methods: Flixweed is a prolific seed-producer, and can build up a reserve of seed in the soil. The seeds survive for years in the soil. Plants are most easily removed when they are small rosettes (ground-hugging forms). Control relies on preventing the production of seed.

Mechanical: Dig, hoe or pull young plants. Use mechanical control methods prior to formation of flowers and seeds. Mow to prevent flowering and production of seed.

Cultural: Plant desirable vegetation to compete with tumble mustard; minimize soil disturbance.

Biological: Do not graze. Eating large quantities of flowers can be toxic to livestock, and can result in death.

Chemical: Apply broadleaf-selective herbicides on young plants.

References:

- Angvick, T. and M. Schat. 2009. HPIPM: Flixweed. Center for Invasive Species and Ecosystem Health, U. of Georgia, <http://wiki.bugwood.org/HPIPM:Flixweed>.
- DiTomaso, J.M. and E.A. Healy. 2007. Weeds of California and Other Western States. University of California Publication #3488.
- Howard, Janet L. 2003. *Descurainia sophia*. In: Fire Effects Information System. U.S.D.A. Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer), <http://www.fs.fed.us/database/feis/plants/forb/dessop/all.html>.
- UC IPM. 2010. Flixweed, <http://www.ipm.ucdavis.edu/PMG/WEEDS/flixweed.html>.
- Whitson, Tom D. (editor). 2002. Weeds of the West. University of Wyoming, Jackson, Wyoming.



The leaves are fern-like in appearance.



Flowers are tiny, yellow and produce long, slender seed pods.



The flower are arranged in a branched structure.

(All photos by S. Donaldson)

The University of Nevada, Reno is an Equal Opportunity/ Affirmative Action employer and does not discriminate on the basis of race, color, religion, sex, age, creed, national origin, veteran status, physical or mental disability, or sexual orientation in any program or activity it conducts. The University of Nevada employs only United States citizens and aliens lawfully authorized to work in the United States.